

Quantum Group based Modeling for the description of high temperature superconductivity and antiferromagnetism: Some supporting evidence

Sher Alam

Physical Science Division, ETL, Tsukuba, Ibaraki 305, Japan

Abstract

Following our recent conjecture to model the phenomena of antiferromagnetism and superconductivity by quantum symmetry groups, we propose in the present note three toy models, namely, one based on $SO_q(3)$ the other two constructed with the $SO_q(4)$ and $SO_q(5)$ quantum groups. Possible motivations and rationale for these choices are outlined. One of the prime motivations underlying our proposal is the experimental observation of *stripe* structure [phase] in high T_c superconductivity [HTSC] materials. A number of experimental techniques have recently observed that the CuO_2 are rather inhomogeneous, providing evidence for phase separation into a two component system. i.e. carrier-rich and carrier-poor regions. In particular, extended x-ray absorption fine structure [EXAFS] demonstrated that these domains forms stripes of undistorted and distorted local structures alternating with mesoscopic length scale comparable with coherence length in HTSC. Evidence for our conjecture is discussed.

The Hubbard Hamiltonian [HH] and its extensions dominate the study of strongly correlated electrons systems and the insulator metal transition [1]. One of the attractive feature of the Hubbard Model is its simplicity. It is well known that in the HH the band electrons interact via a two-body repulsive Coulomb interaction; there are no phonons in this model and neither in general are attractive interactions incorporated. With these points in mind it is not surprising that the HH was mainly used to study magnetism. In contrast superconductivity was understood mainly in light of the BCS theory, namely as an instability of the vacuum [ground-state] arising from effectively attractive interactions between electron and phonons. However Anderson [2,3] suggested that the superconductivity in high T_c material could arise from purely repulsive interaction. The rationale of this suggestion is grounded in the observation that superconductivity in such materials arises from the doping of an otherwise insulating state. Thus following this suggestion the electronic properties in such a high T_c superconductor material close to a insulator-metal transition must be considered. In particular the one-dimensional HH is considered to be the most simple model which can account for the main properties of strongly correlated electron systems including the metal-insulator transition. Long range antiferromagnetic order at half-filling has been reported in the numerical studies of this model [4,5]. Away from half-filling this model has been studied in [6,7].

In theories based on magnetic interactions [2,3] for modelling of HTSC, it has been assumed that the CuO_2 planes in HTSC materials are microscopically homogeneous. However, a number of experimental techniques have recently observed that the CuO_2 are rather inhomogeneous, providing evidence for phase separation into a two component system. i.e. carrier-rich and carrier-poor regions [8]. In particular, extended x-ray absorption fine structure [EXAFS] demonstrated that these domains forms stripes of undistorted and distorted local structures alternating with mesoscopic length scale comparable with coherence length in HTSC. The neutron pair distribution function of Egami et al. [9] also provides structural evidence for two component charge carriers. Other techniques also seem to point that below a certain temperature T^* the CuO_2 planes may have ordered stripes of carrier-rich and carrier-poor domains [9]. The emergence of experimental evidence for inhomogeneous structure has led to renewal of interest, in theories of HTSC which are based on alternative mechanism, such as phonon scattering, the lattice effect on high T_c superconductivity [10–12,9]. Polarized EXAFS study of optimally doped $\text{YBa}_2\text{Cu}_3\text{O}_y$ shows in-plane lattice anomaly [8] below a characteristic temperature $T^{*\dagger}$ which lies above T_c , and close to the characteristic temperature of spin gap opening T^* . It is an interesting question if the in-

The following can be taken as a definition of T^ : T^* is an onset temperature of pseudogap opening in spin or charge excitation spectra.

$\dagger T^{*'} may be defined as follows: $T^{*'}$ is an onset temperature of local phonon anomalies and $T^{*'}$ < T^* .$

plane lattice anomaly is related to the charge stripe or spin-phonon interaction. We note that it has been attempted in [13,14] to relate the spin gap observed in various experiments such as NMR, neutron scattering and transport properties to the short-range ordering of spin singlets.

Zhang [15] proposed a unified theory of superconductivity and antiferromagnetism [‡]based on SO(5) symmetry and suggested that there exists an approximate global SO(5) symmetry in the low temperature phase of the high T_c cuprates. In this model one has a five component order parameter. Three components correspond to a spin one, charge zero particle-hole pair condensed at the center of mass momentum (π, π) , these components represent antiferromagnetic order in the middle of Mott insulating state. The remaining two components correspond to a spin-singlet charge $\pm 2e$ Cooper pair of orbital symmetry $d_{x^2-y^2}$ condensed in zero momentum state, these last two components are supposed to correspond to superconductivity in the doped Mott insulator. Baskaran and Anderson [16] have presented an elegant series of criticisms of the work in Ref. [15]. In view of the critique presented in Ref. [16], and the features of quantum groups we were led to our conjecture that at the simplest level and as a preliminary step, it is tempting to base a model for superconductivity and antiferromagnetism on a quantum group symmetry rather than the usual classical Lie group [26].

We now recall some useful details about quantum groups [19,18] and our simple discussion about quantum groups outlined in [26]. We caution the reader that currently there is no ‘satisfactory’ general definition of a quantum group[§]. However it is commonly accepted [19] that quantum groups are certain ‘well-behaved’ Hopf algebras and that the standard deformations of the enveloping Hopf algebras of semisimple Lie algebras and of coordinate Hopf algebras of the corresponding Lie groups are guiding examples. An amazing feature of quantum group theory is the unexpected connections with apparently unrelated concepts in physics and mathematics such as Lie Groups, Lie Algebras and their representations, special functions, knot theory, low-dimensional topology, operator algebras, noncommutative geometry, combinatorics, quantum inverse scattering problem, theory of integrable models, conformal and quantum field theory and perhaps others.

We consider even the modelling of HTSC materials based on quantum groups as a preliminary step, since it is quite likely that a realistic model which unifies a complex system containing antiferromagnetic and superconducting phases, may require the mathematical machinery currently being used for string theory, or something even beyond it. Another important motivation for our conjecture is to model Stripes. Stripes can be aptly described as being found in the unstable two-phase region between the antiferromagnetic and metal-

[‡]We note that theories of cuprates based upon a quantum critical point have also been suggested by others, see for e.g. Sachdev et al. [17].

[§]We mean in terms of rigorous mathematics

lic states [20]. One of the real challenges is to formulate a theory which gives rise to the equivalent of “Fermi surface”^{**} whose excitation surface derives from *fluctuations that are not uniform in space*. Intuitively one may imagine these non-uniform fluctuations as arising out of a nonlinear sigma-like model. We further conjecture that superconductivity arises when two immiscible phases, namely a 2-D antiferromagnetic state and a 3-D metallic state, are “forced” to meet at $T_c \rightarrow \infty$. As is well-known, ordinary low temperature superconductors arise entirely out of a “metallic” like state. In contrast High T_c superconductors have relatively large T_c since we *cannot* smoothly map two immiscible states [metallic and antiferromagnetic] together. It is the lack of this smooth mapping that is precisely responsible for the High T_c . The lack of smooth mappings may be modelled by the nontrivial phases which arise out of the braiding operations of quantum groups.

In lieu of our remarks on quantum groups [26] to model antiferromagnetism and superconductivity we first consider a simple toy model, namely $SO_q(3)$. Why $SO_q(3)$? To this end we recall the observation that superconductivity in HTSC materials arises from the doping of an otherwise insulating state. On the other hand it is known that the low energy effective Hamiltonian of a antiferromagnet can be described by the $SO(3)$ nonlinear σ model [21]. There are also claims [22–24] that the $SO(3)$ nonlinear sigma model may also suffice to describe the underdoped region of the oxide superconductors. Simply put, experimentally we know that the doping of oxide antiferromagnet leads to superconductivity for a region of the doping parameter. The actual picture is more complicated for example by the presence of stripes. To be concrete we have in mind the generalized phase diagram as seen in La-SrCuO₄ [see Fig. 3.26, page 98 of Ref. [25], where the following phases are enumerated: antiferromagnetic insulator, messy insulating phase, 2D strange metal, 3D metal and superconductor on the temperature versus doping diagram.]. In a simple toy model we may model the antiferromagnetic material plus the doping by $SO_q(3)$, where the quantum group parameter q is related to the physical doping δ . In a naive scenario we may take $\delta = q - 1$, when $q = 1$ $SO_q(3)$ reduces to $SO(3)$, $\delta = 0$ i.e. zero doping and we recover the $SO(3)$ nonlinear σ model description for the low energy effective Hamiltonian.

There are several reasons for choosing $SO_q(4)$ and $SO_q(5)$ As already mentioned $SO(3)$ is a symmetry group for antiferromagnet insulator at the level of effective Hamiltonian. On the other hand effective Hamiltonian of a superconductor may be described by a $U(1)$ nonlinear sigma model [XY model], for example it was indicated in Ref. [27] that the metal insulator transition may be described by the XY model. It was further pointed out in Ref. [28] that superconducting transition on the underdoped side of oxides by a renormalized classical model. In addition it is worth noting that $SO(3)$ spin rotation and $U(1)$ phase/charge rotation are symmetries of the microscopic t-J model. One of the simplest group to embed

^{**}We mean, as is usual [20], that with a metallic state we can always associate a “Fermi surface” which describes the low-energy excitations. This is a finite volume surface in momentum space which arises due to all the one-particle amplitudes [20].

$SO(3) \times U(1)$ is $SO(5)$. Now keeping in mind the discussion in [26] and here we propose to model a unified group for antiferromagnetic, superconducting and other phases in HTSC materials by $SO_q(5)$. Even in the considerations based on $SO(5)$ the concepts of Hopf maps, quaternions, Yang monopole and Berry phase [29] have crept up. This leads more support to our contention that model for HTSC materials must be based on quantum group rather than the classical group in this case $SO_q(5)$ instead of $SO(5)$. A singlet-triplet model has been suggested recently by Mosvkin and Ovchinnikov [30] in light of experimental considerations. This model is based on the Hamiltonian of the two-component spin liquid [30]. The Hamiltonian considered in [30] has $SO(4)$ group symmetry. The calculations and results in [30] are encouraging in that they elaborate and give insight into static and dynamical spin properties of cuprates including paramagnetic susceptibility, nuclear resonance and inelastic magnetic neutron magnetic scattering. However a central issue as recognized in [30] and not dealt with is charge inhomogeneity and phase separation. It is tempting to replace $SO(4)$ by $SO_q(4)$ and see if one could account for charge inhomogeneity and phase separation.

We must also clearly state that we don't at the moment have a magic insight to tell us which specific quantum group must be chosen to model HTSC^{††}. However we feel that the quantum groups arising from the classical orthogonal groups, i.e. $SO(N)$ are a good and worthwhile starting point, since they naturally incorporate the symmetry group of the insulating antiferromagnetic state and are naturally rich enough to accomodate quantum liquid behaviour.

We now give some supporting evidence for our conjecture, in order to establish a link between quantum groups and physics of the high- T_c systems:

1. One of the main theme of quantum groups is the braiding statistics as mentioned previously. In turn one of the consequences of braiding statistics is the existence of non-abelian statistics [see for example [31]]. For example if we choose the group $SO(5)^{\ddagger\ddagger}$ It has been noted in [32] that by perturbative renormalization group analysis two $SO(5)$ invariant fixed points are identified: a massless point, described by $SO(5)_1$ the WZW theory and a massive point corresponding to the Gross-Neveu [GN] field theory. In turn starting from two-leg ladder models subject to $SO(5)$ symmetry one arrives at a qualitative agreement with the above mentioned results based on perturbative renormalization group analysis. The existence of non-abelian electrons can be established in both the critical $SO(5)_1$ WZW theory and GN field theory as noted in [31]. This is not surprising in the quantum group framework where both $SO(5)_1$ WZW theory and GN field theory are special cases [Appendix]. In short if you start with classical

^{††}We have recently proposed $SO(7)$ as a candidate for the classical group underlying the quantum group.

^{‡‡}The choice of a particular group is a separate issue.

SO(5) and apply analysis such as two-leg ladder or perturbative renormalization group you arrive at special cases of corresponding quantum group. Thus our conjecture that one should start with quantum group such as $SO(5)_q$ rather than starting with the classical group SO(5). One can interpret the photoemission experiments as signatures indicating the existence of non-abelian electron statistics which is a natural feature of braiding operations of quantum groups. Thus starting from ladder models obeying classical symmetry naturally leads one to particular cases of quantum groups.

2. Keeping in mind the point above, we propose that we identify fixed points corresponding to various k or q since the two are related and make expansion around these points, to control these expansions we can impose gauge structure and we thus arrive at quantum group based gauge theory or collection of classical gauge theories which represent the condensed matter systems such as cuprates. This is different than ordinary gauge theories in several ways, for in ordinary field theory one has a well-defined critical point here the critical points are not unique or simple. The real transition than in condensed matter system is represented by collection or *ensemble average* of several chosen critical points which come from ordinary field theory [gauge theory] [33].
3. The existence of stripe phase or charge ordering is yet another indication of the quantum group conjecture. As recently emphasized and noted by Emery et al. [34], that stripe phases are predicted and observed in large class of strongly correlated materials, namely doped antiferromagnets [such as copper oxide superconductors]. Prominent in understanding the stripe formation is the mechanism of dynamical dimensional reduction, which simply means that a synthetic metal will behave over a wide range of temperature and energy as if it were of *lower dimensionality*. *As is well-known one place quantum groups arise is out of symmetries of one dimensional systems*. And they are connected to existence of non-trivial topology [braiding] in two dimensions. The dimensional reduction to us is an evidence for our conjecture. In other words the existence of stripes is the indication of the underlying quantum group symmetry.
4. Another recent claim [36] if true^{§§} further supports our intuition. The fluctuations associated with a striped phase are expected to be one-dimensional, whereas the magnetic fluctuations display two dimensional symmetry. It is claimed in [36] that this apparent two-dimensionality results from measurements on twinned crystals, and that similar measurements on substantial de-twinned crystals of $YBa_2Cu_3O_{6.6}$ reveal the one-dimensional character of the magnetic fluctuations thus making the stripe scenario more stronger. And as noted above quantum groups are related to the symmetries of one-dimensional systems.

^{§§}This is especially encouraging since we did not know about the one-dimensional nature of magnetic fluctuations when we made our conjecture.

5. Experimental evidence taken together about charged and magnetic stripes indicates that the superconducting state is influenced by one-dimensional fluctuations which are intimately tied to stripes. There is an intimate connection between sigma models, Kac-Moody algebras, quantum algebra and rich structure of strings. The noncommutativity which is intimately tied to quantum groups can be a powerful tools to estimate the effects of the one-dimensional fluctuations [charge, magnetic and lattice] on the superconducting state.

One expects that the various phases, namely electronic liquid crystal states can be classed under quantum group symmetries. The quantum Hall systems indicate this trend.

In conclusion, the quantum orthogonal groups $SO_q(N)$ are proposed as potential candidate for modelling the theory of HTSC materials. A strong feature of quantum groups is that they unify classical Lie algebras and topology. In more general sense it is expected that quantum groups will lead to a deeper understanding of the concept of symmetry in physics.

ACKNOWLEDGMENTS

The author's work in this paper is supported by the Japan Society for the Promotion of Science [JSPS].

APPENDIX:

We recall [18,26] that

$$q \leftrightarrow e^{2\pi i/(k+2)} \tag{A1}$$

If we make the above correspondence it can be shown by examining various identities of WZW model that the braiding properties of WZW model at level k are determined by the representation theory of quantum groups. As a trivial check if one sets $q = 1$ in A1, which is the limit in which quantum group reduces to the ordinary classical group, then the right-hand side of A1 we must set $k \rightarrow \infty$, which is precisely the limit in which Kac-Moody algebra reduces to ordinary classical algebra. We recall that the symmetry generators of the WZW model obey a special case of Kac-Moody algebra.

Some remarks about classical groups, sigma models, and relations of non linear sigma models to quantum groups and strings are included in [33,35], for more details see kak91.

REFERENCES

- [1] E. Fradkin, *Field Theories of Condensed Matter Systems*, Addison-Wesley Publishing Company, 1991.
- [2] P. W. Anderson, *Science* **235**, (1987), 1196.
- [3] P. W. Anderson et al., *Phys. Rev. Lett.* **58**, (1987), 2790.
- [4] J. Hirsch and S. Tang, *Phys. Rev. Lett.* **62**, (1989), 591.
- [5] A. P. Balachandran et al., *Hubbard Model and Anyon Superconductivity*, World Scientific, 1990.
- [6] A. Moreo et al., *Phys. Rev.* **B41**, (1990), 2313.
- [7] A. Moreo and D. J. Scalapino, *Phys. Rev.* **B43**, (1991), 8211.
- [8] H. Oyanagi et al., ‘Local Lattice Distortions in $\text{YBa}_2\text{Cu}_3\text{O}_y$: Doping dependence, in International Conference on Stripes in High T_c Superconductivity, Rome, June, 1998. Proceedings of STRIPES98, to appear in *J. Superconductivity*, (1999).
- [9] T. Egami and S. J. L. Billinge, *Prog. Mater. Sci* **38**, (1994), 359.
- [10] *Lattice Effects in High T_c Superconductors*, ed by Y. Bar-Yam et al., World Scientific Publishing, Singapore, 1992.
- [11] *Phase Separation in Cuprate Superconductors*, ed by K. A. Muller and G. Benedek, World Scientific Publishing, Singapore, 1993.
- [12] *Phase Separation in Cuprate Superconductors*, ed by E. Sigmund and K. A. Muller, World Scientific Publishing, Singapore, 1994.
- [13] T. Tanamoto et al., *J. Phys. Soc. Jpn.* **61**, (1992), 1886.
- [14] T. Tanamoto et al., *J. Phys. Soc. Jpn.* **62**, (1993), 717.
- [15] Shou-Cheng Zhang, *Science* **275**, (1997), 1089.
- [16] G. Baskaran and P. W. Anderson, cond-mat/9706076.
- [17] S. Sachdev et al., *Phys. Rev.* **B51**, (1995), 14874.
- [18] M. Kaku, *Strings, Conformal Fields, and Topology: An Introduction*, Springer-Verlag, 1991.
- [19] A. Klimyk and K. Schmüdgen, *Quantum Groups and their Representations*, Springer-Verlag, 1997.
- [20] P. W. Anderson and G. Baskaran, cond-mat/9711197.
- [21] S. Chakravarty et al., *Phys. Rev.* **B39**, (1994), 2344.
- [22] D. Pines, *Physica* **C235**, (1994), 113.
- [23] A. V. Barzykin and D. Pines, *Phys. Rev.* **B52**, (1995), 13585.
- [24] A. V. Chubukov et al., *Phys. Rev.* **B49**, (1994), 11919.
- [25] P. W. Anderson, *The Theory of Superconductivity in the High- T_c Cuprates*. (Princeton University Press, Princeton, New Jersey 1999).
- [26] S. Alam, *A Conjecture for possible theory for the description of high temperature superconductivity and antiferromagnetism*. Proceedings of Quantum Phenomena in Advanced Materials at High Magnetic Fields, 4th International Symposium on Advanced Physical Fields [APF-4]. KEK-TH-607, KEK Preprint 98-xxx, cond-mat/9812060.
- [27] S. Doniach and M. Inui, *Phys. Rev.* **B41**, (1990), 6668.

- [28] V. J. Emery and S. Kivelson, *Nature* **374**, (1995), 434.
- [29] Shou-Cheng Zhang, cond-mat/9808309.
- [30] A. S. Moskvina and A. S. Ovchinnikov, *Physica C* **296**, (1998), 250.
- [31] P. Bouwknecht and K. Schoutens, *Phys. Rev. Lett.* **82**, (1999), 2757.
- [32] H. -H. Lin, *Phys. Rev.* **B58**, (1998), 1794.
- [33] Sher Alam et al., *The patching of critical points using quantum groups*, cond-mat/0004350.
- [34] V. J. Emery et al., cond-mat/9907228.
- [35] Sher Alam et al., *The Choice of symmetry group for cuprates*, cond-mat/0004269.
- [36] H. A. Mook et al., *Nature* **404**, (2000), cond-mat/0004362.